


Epigallocatechin 3-O-Gallate Extract from Green Tea

Application
Food Analysis
Robert Ricker

Epigallocatechin 3-O-gallate (EGCG) belongs to a class of compounds called flavonoids and is further subclassified as a flavanol, due to the level of oxidation in its chemical structure. This compound has been recognized as a cancer-preventive compound due to its ability to inhibit urokinase, an enzyme which has been associated with accelerated tumor cell growth. Due to the interest in holistic-type, preventative medicine approaches in today's society, a method was developed for a series of catechins and an actual extract of green tea to serve as a general interest application.

Highlights

- Good peak shape and resolution are maintained for a group of catechins on Agilent ZORBAX SB-C8 at low pH.
- Sterically protected bonded phases, like SB-C8, offer extended column lifetime even with TFA-containing (low-pH) mobile phases.
- Good retention of the catechins allows adequate separation from other UV-absorbing compounds in the actual tea extract.



ZORBAX SB-C8 (4.6 x 150 mm; 3.5 μ m) (Agilent Part No. 863953-906)
Mobile Phase: 75% 0.1% Trifluoroacetic acid: 25% Methanol
Inj. Vol. 5 μ L, 1 mL/min, 40°C
Det. UV (280 nm)



Agilent Technologies

*Robert Ricker is an application chemist
based at Agilent Technologies, Wilmington,
Delaware.*

For more information on our products and
services, visit our website at:
www.agilent.com/chem

Copyright© 2002 Agilent Technologies, Inc.
All Rights Reserved. Reproduction,
adaptation or translation without prior
written permission is prohibited, except as
allowed under the copyright laws.

Agilent shall not be liable for errors
contained herein or for incidental or
consequential damages in connection with
the furnishing, performance, or use of this
material.

Information, descriptions, and specifications
in this publication are subject to change
without notice.

Printed in the USA
April 25, 2002
5988-6357EN



Agilent Technologies